




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07F 15/00, G01N 33/58, 33/72, C07K 5/10, 14/805</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/62918 (43) International Publication Date: 9 December 1999 (09.12.99)</p>
<p>(21) International Application Number: PCT/US99/11855 (22) International Filing Date: 28 May 1999 (28.05.99) (30) Priority Data: 60/087,576 1 June 1998 (01.06.98) US (71) Applicant: ROCHE DIAGNOSTICS CORPORATION [US/US]; 9115 Hague Road, Indianapolis, IN 46250 (US). (72) Inventors: BUCK, Harvey, B., Jr.; 8147 Bay Brook Drive, Indianapolis, IN 46256 (US). DENG, Zhe, D.; 13722 Goldfinch Drive, Carmel, IN 46032 (US). (74) Agent: LAMMERT, Steven, R.; Barnes & Thornburg, 11 South Meridian Street, Indianapolis, IN 46204 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>

(54) Title: REDOX REVERSIBLE IMIDAZOLE-OSMIUM COMPLEX CONJUGATES



(57) Abstract

Novel imidazole-osmium complex conjugates, their preparation, and their use in electrochemical assays are described. The redox reversible-osmium complexes can be prepared to exhibit unique reversible redox potentials and can thus be used in combination with other electroactive redox reversible species having redox potentials differing by at least 50 millivolts in electrochemical assays designed for use of multiple electroactive species in the same cell and in the same sample without interference between the two or more redox coupled conjugate systems.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

REDOX REVERSIBLE IMIDAZOLE
-OSMIUM COMPLEX CONJUGATES

5 Field of the Invention

 This invention relates to novel redox-reversible conjugates. More particularly the invention is directed imidazole complexed-osmium conjugates useful for detection and quantification of biologically significant analytes in a liquid sample.

10 Background and Summary of the Invention

 Therapeutic protocols used today by medical practitioners in treatment of their patient population requires accurate and convenient methods of monitoring patient disease states. Much effort has been directed to research and development of methods for measuring the presence and/or concentration of biologically significant
15 substances indicative of a clinical condition or disease state, particularly in body fluids such as blood, urine or saliva. Such methods have been developed to detect the existence or severity of a wide variety of disease states such as diabetes, metabolic disorders, hormonal disorders, and for monitoring the presence and/or concentration of ethical or illegal drugs. More recently there have been significant advancements in the
20 use of affinity-based electrochemical detection/measurement techniques which rely, at least in part, on the formation of a complex between the chemical species being assayed (the "analyte") and another species to which it will bind specifically (a "specific binding partner"). Such methods typically employ a labeled ligand analog of the target analyte, the ligand analog selected so that it binds competitively with the
25 analyte to the specific binding partner. The ligand analog is labeled so that the extent of binding of the labeled ligand analog with the specific binding partner can be measured and correlated with the presence and/or concentration of the target analyte in the biological sample.

 Numerous labels have been employed in such affinity based sample
30 analysis techniques, including enzyme labeling, radioisotopic labeling, fluorescent labeling, and labeling with chemical species subject to electrochemical oxidation and/or reduction. The use of redox reversible species, sometimes referred to as electron transfer agents or electron mediators as labels for ligand analogs, have proven to provide a practical and dependable results in affinity-based electrochemical assays.

However, the use of electrochemical techniques in detecting and quantifying concentrations of such redox reversible species (correlating with analyte concentrations) is not without problem. Electrochemical measurements are subject to many influences that affect the accuracy of the measurements, including not only those relating to variations in the electrode structure itself and/or matrix effects deriving from variability in liquid samples, but as well those deriving from interference between multiple electroactive species, especially when assay protocols require detection or quantification of multiple electroactive species.

The present invention relates to novel diffusible, redox-reversible osmium-imidazole conjugates useful in immunosensors based on either indirect amplified electrochemical detection techniques or on direct electrochemical measurement of detectable species with microarray electrodes under bipotentiostatic control. An Os-imidazole complex can, for example be covalently attached to a peptide which has amino acid sequence of the binding epitope for an antibody. When Os complex/peptide conjugate is bound to antibody, the conjugate does not function electrochemically; it is said to be "inhibited". Typically an analyte present in sample will compete with Os-imidazole complex/peptide conjugate for the limited number of binding sites on the antibody. When more analyte is present, more free Os-imidazole complex/peptide conjugate will be left in an unbound diffusible state producing higher current at a sensor electrode, *i.e.* one of the working electrodes where measured events (oxidation or reduction) are taking place. In the opposite case, when less analyte is present, more indicator/peptide conjugate will be bound to antibody resulting less free conjugates and producing lower current levels at the working electrodes. Therefore the current detected at either one of the working electrodes will be a function of analyte concentration.

It is frequently desired to measure more than one analyte species in a liquid sample. Measurement of multiple species in a mixture has been achieved with photometry and fluorescence, via selection of the appropriate wavelengths. Electrochemical measurements of a single species in a complex mixture are routinely made by selecting a potential at which only the desired species is oxidized or reduced (amperometry) or by stepping or varying the potential over a range in which only the desired species changes its electrochemical properties (AC and pulse methods). These

methods suffer from disadvantages including lack of sensitivity and lack of specificity, interference by charging and matrix polarization currents (pulse methods) and electrode fouling due to the inability to apply an adequate over potential. Moreover, electrochemical measurements are complicated by interference between the multiplicity
5 of electroactive species commonly extant in biological samples.

Electrode structures which generate steady state current via diffusional feedback, including interdigitated array electrodes (IDAs) (Figs. 1 and 2) and parallel plate arrangements with bipotentiostatic control are known. They have been used to measure reversible species based on the steady state current achieved by cycling of the
10 reversible species. A reversible mediator (redox reversible species) is alternately oxidized and reduced on the interdigitated electrode fingers. The steady state current is proportionate to mediator concentration (Fig. 3) and limited by mediator diffusion. A steady state current is achieved within seconds of applying the predetermined anodic (more positive) and cathodic (less positive or negative) potentials (Fig. 6) to the
15 microelectrode array. The slope of a plot of the IDA current vs. mediator concentration is dependent on IDA dimensions, and the slope increases with narrower electrode spacings (Fig. 7).

The present invention provides novel osmium-imidazole complex conjugates useful in a method for measuring multiple analyte species in the same
20 sample, and optimally on the same electrode structure, thus improving the accuracy of the relative measurements. The present conjugates can be used with other electroactive conjugate species having unique redox potentials to provide an electrochemical biosensor with capacity to provide improved accuracy. Analyte concentration can be measured/calculated from electrometric data obtained on the
25 same liquid sample with the same electrode structure, thereby minimizing perturbations due to variability in sample or electrode structure.

The diffusible osmium conjugates of this invention find use in assays based on the principle of diffusional recycling, where a diffusible redox reversible species is alternately oxidized and reduced at nearby electrodes (the working
30 electrodes), thereby generating a measurable current. As alternate oxidation and reduction is required for measurement, only electroactive species which are electrochemically reversible at the predetermined redox potential are measured thereby

eliminating, or at least reducing, the impact or interference from non-reversible electroactive species in the sample or other reversible-redox species having unique (at least 50 millivolts different) redox potential. Redox reversible species having different oxidation potentials can be independently measured in a mixture by selecting and
5 bipotentially controlling the oxidizing and reducing potentials for neighboring electrode pairs so that only the species of interest is oxidized at the anode and reduced at the cathode. When the working electrodes are dimensioned to allow diffusional recycling of the redox-reversible-species at the selected oxidizing and reducing potentials appropriate for that species, a steady state current is quickly established
10 through the sample and the electrode structure. The magnitude of the current at the working electrodes where the measurable oxidative and reductive events are taking place, is proportional to the concentration of the diffusible redox reversible species in the sample. When two or more redox reversible species are utilized, they are selected to have redox potentials differing by at least 50 millivolts, most preferably at least 200
15 millivolts, to minimize interference between one species and the other in measurements of the respective steady state currents. The present osmium complex conjugates have unique redox potentials that allow them to be used with/in the presence of other electroactive conjugates without (or with minimal) interference.

The present conjugates can be used in any electrode structure/system
20 which allows for diffusional recycling to achieve steady state current in response to application of pre-selected complex species-specific anodic and cathodic potentials. Suitable electrode structures include interdigitated array microelectrodes and parallel plate electrodes separated by distances within the diffusion distance of the respective redox reversible species. The electrode structures typically include a reference
25 electrode (*e.g.*, Ag/AgCl), at least two working electrodes, and optionally an auxiliary electrode for current control. In use, a programmable bipotentiostat is placed in electrical communication with the electrode structure for applying the respective anodic and cathodic potentials specific for each of the respective redox reversible species utilized in the method/biosensor. Several novel osmium complexes including
30 those of this invention have been developed for use as labels for preparing ligand analog conjugates having potential differences sufficient (at least 50 millivolts) to allow

the use of two osmium complexes (as opposed to an osmium complex and a ferrocene or other redox reversible label) in multiple conjugate based electrochemical assays.

The present osmium conjugates are useful in a method for measuring the concentration of one or more analytes in a liquid sample. The method includes

5 contacting a portion of the sample with pre-determined amounts of at least a first and second redox reversible species having a redox potential differing by at least 50 millivolts from that of each other species. Each respective species comprises a liquid sample diffusible conjugate of a ligand analog of an analyte in the liquid sample and a redox reversible label. The liquid sample is also contacted with a predetermined

10 amount of at least one specific binding partner for each analyte to be measured. The diffusible conjugate is selected so that it is capable of competitive binding with the specific binding partner for said analyte. The concentration of the diffusible redox-reversible-species in the liquid sample is then determined electrochemically. The sample is contacted with an electrode structure, including a reference electrode and at

15 least first and second working electrodes dimensioned to allow diffusional recycling at least one of the diffusible redox-reversible-species in the sample, when a predetermined redox-reversible-species-dependent cathodic potential is applied to one working electrode and a predetermined redox-reversible-species-dependent anodic potential is applied to the second working electrode. Typically, a first cathodic potential is applied

20 to the first working electrode and a first anodic potential is applied to the second working electrode to establish current flow through the sample due to diffusional recycling of the first redox-reversible-species without significant interference from the second redox-reversible-species. Current flow through one or more of the electrodes at the first anodic and cathodic potentials is measured. Similarly current flow

25 responsive to application of second cathodic and anodic potentials to electrodes in contact with the sample is measured and correlated with measured current flows for known concentrations of the respective redox-reversible-species, said concentrations being proportionate to the respective analyte concentrations.

The reagent components including the present imidazole-osmium

30 conjugates of the invention and the specific binding partners, can be provided in the form of a test kit for measuring the targeted analyte(s) in a liquid sample, either as separate reagents or, more preferably, combined as a multi-reagent composition, e.g.

combined redox reversible species, combined specific binding partners, or combined redox reversible species and specific binding partners. The kit optionally, but preferably, includes an electrode structure dimensioned to allow diffusional redox recycling of diffusible redox reversible species in the liquid sample. The electrode
5 structure includes conductors for connecting the structure to a bipotentiostat programmed to apply redox-reversible-species-dependent-anodic and cathodic potentials to the electrode structure and to sense and measure current flow, typically at one or both of the working electrodes, responsive to such potentials.

This invention is based on the preparation and use of novel
10 electrochemically detectable osmium complexes and covalent conjugates of said complexes having oxidation potentials differing sufficiently from other redox-reversible complexes to enable their use together with other osmium or other metal conjugates. Thus, there are provided novel osmium labeled ligand analogs capable of binding to a specific binding partner of a biologically significant analyte. The electrochemically
15 detectable osmium conjugates comprise a *bis*(bipyridyl) imidazolyl haloosmium complex characterized by fast mediation kinetics and low redox potential (+150mV vs. Ag/AgCl). Another group of osmium complex labeled, electrochemically detectable conjugates that can be used with the present complexes in multi-conjugate assay protocols include *tris*(bipyridyl) osmium complexes, which, like the *bis*(bipyridyl)
20 imidazolyl haloosmium complexes are characterized by fast mediation kinetics, but the *tris*(bipyridyl) complexes have a redox potential sufficiently different from the *bis*(pyridyl) imidazolyl chloroosmium complexes to allow their use together in assays utilizing microelectrode arrays for measuring more than one analyte in a single liquid sample by concentration dependent currents amplified by diffusional redox recycling.

25 The present osmium complex conjugates can be used in combination with another conjugated redox-reversible-species for the measurement of both glycosylated hemoglobin and hemoglobin in a lysed blood sample. One redox-reversible-species preferably comprises an imidazole-osmium complex covalently linked to a ligand analog of either hemoglobin or glycosylated hemoglobin, and a
30 second redox-reversible-species comprising a second redox reversible label covalently bound to a ligand analog of the other of the two target analytes. The method enables measurement of the concentration of both the glycosylated hemoglobin (HbA1c) and

the concentration of either total hemoglobin or that of unglycosylated hemoglobin (HbA₀) thereby enabling calculation of the results as a ratio of the two measurements (% HbA_{1c}). It is advantageous to assay both HbA_{1c} and total hemoglobin (or HbA₀) using the same principle in a single sample, particularly because ratioing works
5 minimize biases due to environmental effects.

Brief Description of the Drawings

Fig. 1 is an enlarged plan view of an interdigitated array electrode for reversible mediator measurement.

10 Fig. 2 is a partial cross-sectional view of the electrode of Fig. 1 illustrating the conditions of steady state current limited by diffusion of mediator (M).

Fig. 3 is a graphic presentation of dose response currents for a *bis*-(bipyridyl) imidazolyl chloroosmonium mediator peptide conjugate of this invention.

15 Fig. 4 is a graphic illustration of current flow vs. concentration of glycosylated hemoglobin (HbA_{1c}) in blood samples using an osmium conjugate of this invention and enzyme amplified DC amperometry.

Fig. 5 is a graphic illustration of the inhibition of current flow due to free conjugate as a function of antibody concentration (C_n) as measured using enzyme amplified DC amperometry [C₁, >C₂, >C₃].

20 Fig. 6 is a graphic illustration of current flow vs. time using an interdigitated array electrode.

Fig. 7 is a graphic illustration of the effect of the dimensions of the interdigitated array electrode structure on current flow as a function of concentration of an osmium conjugate (Os-DSG-A1c).

25 Fig. 8 is a graphic illustration of current flow as a function of applied potential for a liquid sample containing equimolar (50μM) of a *bis*-(bipyridyl) imidazolyl chloroosmium complex of this invention and a *tris*(bipyridyl) osmium complex.

30 Fig. 9 is a graphic presentation of current flow vs. concentration of a ferrocene-biotin conjugate in the presence of varying amounts of an osmium complex conjugate on interdigitated array electrodes with bipotentiostatic control.

Fig. 10 is a graphic illustration of the effect of concentration of an unlabeled conjugate (BSA-A1c) on current flow in a solution containing osmium labeled conjugate (osmium-DSG-A1c) in the presence of three separate A1c-recognizing antibody compositions.


Fig. 11 illustrates the structure of a *tris*(bipyridyl) osmium labeled conjugate.

Figs. 12-14 are similar and each depict the chemical structure of a *bis*(bipyridyl) imidazolyl chlorosmium labeled peptide conjugate in accordance with this invention.

Detailed Description of the Invention

The diffusible redox reversible species of this invention is a liquid-sample-diffusible conjugate of a ligand analog of an analyte and a redox reversible imidazole-osmium complex. The term "redox reversible" as used herein refers to a chemical species capable of reversible oxidation and reduction in a liquid sample. Redox reversible labels are well-known in the art and include ligand complexes of transition metal ions, for example iron (ferrocene and ferrocene derivatives), ruthenium and osmium. The conjugate is prepared by linking the ligand analog to the label either covalently through difunctional linking agents or by combination of covalent linkages and art-recognized specific binding entities (for example, biotin-avidin).

More particularly, the present invention is directed to a redox reversible osmium complex of the formula




wherein

R and R₁ are the same or different and are 2,2'-bipyridyl, 4,4'-disubstituted-2,2'-bipyridyl, 5-5'-disubstituted-2,2'-bipyridyl, 1,10-phenanthroline, 4,7-disubstituted-1,10-phenanthroline, or 5,6-disubstituted-1,10-phenanthroline, wherein each substituent is a methyl, ethyl, or phenyl group,

- 5 R and R₁ are coordinated to Os through their nitrogen atoms;
 q is 1 or 0;
 R₇ is B-(L)_k-Q(CH₂)_i-;
 R₂ is hydrogen, methyl, or ethyl when q is 1, and R₂ is B-(L)_k-Q(CH₂)_i-
 when q is 0; wherein in the group B-(L)_k-Q(CH₂)_i-
 10 Q is O, S, or NR₄ wherein R₄ is hydrogen, methyl or ethyl;
 -L- is a divalent linker;
 k is 1 or 0;
 i is 1, 2, 3, 4, 5 or 6; and
 B is hydrogen or a group comprising a ligand capable of binding
 15 to a specific binding partner;
 Z is chloro or bromo;
 m is +1 or +2;
 X is monovalent anion, e.g., chloride, bromide, iodide, fluoride,
 tetrafluoroborate, perchlorate, nitrate, or a divalent anion, e.g., sulfate, carbonate, or
 20 sulfite;
 Y is monovalent anion, e.g., chloride, bromide, iodide, fluoride,
 tetrafluoroborate, perchlorate or nitrate; and
 n is 1 or zero,
 provided that when X is a divalent anion, n is zero,
 25 and when m is 1, n is zero and X is not a divalent anion.

Another redox reversible osmium complex that can be used with the present imidazole-osmium conjugates is a compound of the formula



10 wherein

R and R₁ are the same or different and are 2,2'-bipyridyl, 4,4'-disubstituted-2,2'-bipyridyl, 5-5'-disubstituted-2,2'-bipyridyl, 1,10-phenanthrolinyl, 4,7-disubstituted-1,10-phenanthrolinyl, or 5,6-disubstituted-1,10-phenanthrolinyl, wherein each substituent is a methyl, ethyl, or phenyl group,

15 R₅ is 4-substituted-2,2'-bipyridyl or 4,4'-disubstituted-2,2'-bipyridyl wherein the substituent is the group B-(L)_k-Q(CH₂)_i- and the 4'-substituent is a methyl, ethyl or phenyl group;

R, R₁ and R₅ are coordinated to Os through their nitrogen atoms;

Q is O, S, or NR₄ wherein R₄ is hydrogen, methyl or ethyl;

20 -L- is a divalent linker;

k is 1 or 0;

i is 1, 2, 3, 4, 5 or 6;

B is hydrogen or a group comprising a ligand capable of binding to a specific binding partner;

25 d is +2 or +3;

X and Y are anions selected from monovalent anions, chloride, bromide, iodide, fluoride, tetrafluoroborate, perchlorate, and nitrate and divalent anions, e.g., sulfate, carbonate or sulfite wherein x and y are independently 0, 1, 2, or 3 so that the net charge of X_xY_y is -2 or -3.

30 Redox reversible conjugate species of each of those formulas are prepared from the corresponding compounds wherein k is 0 and B is hydrogen by reacting such compounds with either a heterofunctional crosslinker of the formula S-

L'-T wherein L' is a divalent linker and S and T are different electrophilic groups capable of reacting with a nucleophilic group to form a covalent bond, or with a homofunctional crosslinker of the formula S-L'-T wherein L' is a divalent linker and S and T are the same electrophilic groups capable of reacting with a nucleophilic group to form a covalent bond. The resulting products are then reacted with ligand analogs using classical coupling reaction conditions to product the conjugate species. The oxidizing potentials of the respective *bis*(bipyridyl) and *tris*(bipyridyl) osmium complexes defined above is such that the respective complexes can be used as reversible redox labels for the respective redox reversible species in performance of the method. Fig. 8 illustrates a cyclic voltammogram for a liquid sample containing equimolar (50 μ M) amounts of a *bis*(bipyridyl) imidazolyl chloroosmium complex and a *tris*(bipyridyl) osmium complex.

In one embodiment of the invention the specific binding partner for each analyte is an antibody and the ligand analog is selected so that it binds competitively with the analyte to the antibody. There are, however, other examples of ligand-specific binding partner interactions that can be utilized in developing applications of the present method. Examples of ligands and specific binding partners for said ligands are listed below.

20	<u>Ligand</u>	<u>Specific Binding Partner</u>
	Antigen (e.g., a drug substance)	Specific antibody
	Antibody	Antigen
	Hormone	Hormone receptor
25	Hormone receptor	Hormone
	Polynucleotide	Complementary polynucleotide strand
	Avidin	Biotin
	Biotin	Avidin
	Protein A	Immunoglobulin
30	Immunoglobulin	Protein A
	Enzyme	Enzyme cofactor (substrate)
	Enzyme cofactor (substrate)	Enzyme
	Lectins	Specific carbohydrate
35	Specific carbohydrate of lectins	Lectins

The term "antibody" refers to (a) any of the various classes or subclasses of immunoglobulin, *e.g.*, IgG, IgM, derived from any of the animals conventionally used, *e.g.*, sheep, rabbits, goats or mice; (b) monoclonal antibodies; (c) intact molecules or "fragments" of antibodies, monoclonal or polyclonal, the fragments being those which contain the binding region of the antibody, *i.e.*, fragments devoid of the Fc portion (*e.g.*, Fab, Fab¹, F(ab')₂) or the so-called "half-molecule" fragments obtained by reductive cleavage of the disulfide bonds connecting the heavy chain components in the intact antibody. The preparation of such antibodies are well-known in the art.

10 The term "antigen" used in describing and defining the present invention includes both permanently antigenic species (for example, proteins, peptides, bacteria, bacteria fragments, cells, cell fragments and viruses) and haptans which may be rendered antigenic under suitable conditions.

15 The present osmium labeled conjugates are useful alone or in combination with other conjugates in methods for measuring the concentration of one or more analytes in a liquid sample. One method enables two or more independent amperometric measurements of the sample on a single electrode structure. The method comprises

 contacting a volume of said liquid sample with

20 1) predetermined amounts of at least a first and second redox reversible species, each respective species having a redox potential differing by at least 50 millivolts from that of each other species, at least one species comprising a liquid sample diffusible conjugate of a ligand analog of an analyte in the liquid sample and a redox reversible label, said conjugate capable of competitive binding with a specific binding partner for said analyte, and

25 2) a predetermined amount of at least one specific binding partner for each analyte to be measured; and

 electrochemically determining the concentration of each of said diffusible redox-reversible species in the liquid sample by

30 contacting said sample with an electrode structure including a reference electrode and at least first and second working electrodes dimensioned to allow diffusional recycling of the diffusible redox reversible species in the sample when

a predetermine redox-reversible-species-dependent cathodic potential is applied to one working electrode and a predetermined redox-reversible-species-dependent anodic potential is applied to a second working electrode, said diffusional recycling of said species being sufficient to sustain a measurable current through said sample,

5 applying a first cathodic potential to the first working electrode and a first anodic potential to the second working electrode, said first cathodic and anodic potentials corresponding to those respective potentials necessary to establish current flow through the sample due to diffusional recycling of the first redox reversible species without significant interference from said second redox reversible species,

10 measuring current flow at said first anodic and cathodic potentials,

 applying a second cathodic potential to said first or second working electrode and a second anodic potential to the other working electrode, said

15 second cathodic and anodic potential corresponding to those respective potentials necessary to establish current flow through the sample due to diffusional recycling of the second redox-reversible-species without significant interference from the first redox reversible species,

 measuring current flow at said second anodic and cathodic

20 potentials, and

 correlating the respective measured current flows to that for known concentrations of the respective diffusible redox reversible species, said concentrations being proportionate to the respective analyte concentrations.

That method has very broad applicability but in particular may be used to assay: drugs, hormones, including peptide hormones (*e.g.*, thyroid stimulating hormone (TSH), luteinizing hormone (LH), follicle stimulating hormone (FSH), insulin and prolactin) or non-peptide hormones (*e.g.*, steroid hormones such as cortisol, estradiol, progesterone and testosterone, or thyroid hormones such as thyroxine (T4) and triiodothyronine), proteins (*e.g.*, human chorionic gonadotropin (hCG), carcino-embryonic antigen (CEA) and alphafetoprotein (AFP)), drugs (*e.g.*, digoxin), sugars, toxins or vitamins.

The method can be performed on liquid samples comprising biological fluids such as saliva, urine, or blood, or the liquid sample can be derived from environmental sources. The liquid samples can be analyzed "as is," or they can be diluted, buffered or otherwise processed to optimize detection of the targeted
5 analyte(s). Thus, for example, blood samples can be lysed and/or otherwise denatured to solubilize cellular components.

The method can be performed using widely variant sampling handling techniques. Thus, the sample can be premixed with either or both of the specific binding partner for the targeted analytes and the redox reversible species prior to
10 contacting the sample with the electrode structure, or the liquid sample, either neat or pre-processed, can be delivered to a vessel containing predetermined amounts of the redox reversible species and the specific binding partner for subsequent or simultaneous contact with the electrode structure. The order of introduction of the components into the sample is not critical; however, in one embodiment of the
15 invention the predetermined amounts of the specific binding partners are first added to the sample, and thereafter, there is added the predetermined amounts of the redox reversible species. It is also possible to combine the predetermined amounts of the specific binding partners with the redox reversible species to form the respective complexes prior to combining those components with the liquid sample. In that latter
20 case the redox reversible species will be displaced from its respective specific binding partner by the corresponding analyte to provide a concentration of the redox reversible species proportionate to the concentration of analyte in the liquid sample. The reagents, that is, the predetermined amounts of the specific binding partner of each analyte and the predetermined amounts of the corresponding redox reversible species
25 can, for example, be deposited in a vessel for receiving a predetermined volume of the liquid sample. The liquid sample is added to the vessel, and thereafter, or simultaneously, the liquid sample is contacted with the electrode structure.

The electrode structure includes a reference electrode and at least first and second working electrodes dimensioned to allow diffusional recycling of the
30 diffusible redox reversible species in the sample when predetermined redox-reversible-species-dependent-cathodic and anodic potential is applied to the working electrodes. The term "working electrode" as used herein refers to an electrode where measured

events (*i.e.* oxidation and/or reduction) take place and resultant current flow can be measured as an indicator of analyte concentration. "Anodic potential" refers to the more positive potential (applied to the anode) and "cathodic potential" refers to the less positive or negative potential applied to the cathode (vs a reference electrode).

5 Electrodes dimension to allow diffusional recycling are well known in the art and are typically in the form of arrays of microdiscs, microholes, or microbands. In one embodiment the electrodes are in the form of an interdigitated arrangement of microband electrodes with micron or submicron spacing. Short average diffusional length and a large number of electrodes are desirable for effective current amplification
10 by recycling of reversible redox species. The microelectrode arrays can be fabricated, for example, as pairs of interdigitated thin film metal electrodes in micron and submicron geometry arranged on an insulator substrate, for example, oxidized silicon. Each of the electrode fingers (Fig. 1) are spaced from its neighboring finger in the nanometer to low micrometer (1-10 microns) range. Microelectrode arrays can be
15 fabricated using photolithography, electron beam lithography, and so-called lift-off technique. Thus, an interdigitated electrode array (IDA) can be deposited on glass, silicon or polyamide utilizing the following general procedure:

1. Grow thermal oxide layer on silicon substrate;
2. Sputter 400Å chromium seed layer, 2000 Å gold;
- 20 3. Spin-coat and soft-bake photo resist;
4. Expose and develop photo resist with IDA pattern;
5. Pattern gold and chromium with ion beam milling;
6. Strip photo resist; and
7. Cut electrodes into chips by first coating with a protective layer,
25 cutting into strips, stripping the protective layer, and cleaning electrode surfaces in oxygen plasma.

The electrode structure can be formed on an inner surface of a chamber for receiving the liquid sample, *e.g.*, a cuvette, a capillary fill chamber, or other sample receiving vessel wherein the electrode structure can be contacted with the liquid
30 sample. Alternatively, the electrode structure can form part of a probe for dipping into the liquid sample after the sample has been contacted with the predetermined amounts of the redox reversible species and the specific binding partners. The electrode

structure is in contact with conductors that enable application of the respective cathodic and anodic potentials for carrying out the present method. The anodic and cathodic potentials are applied relative to a reference electrode component of the electrode structure using a bipotentiostat. The electrode structure can optionally
5 include an auxiliary electrode for current control. The bipotentiostat is utilized to apply a first cathodic potential to a first working electrode and a first anodic potential to a second working electrode, the first cathodic and anodic potentials corresponding to those respective potentials necessary to establish current flow through the sample due to diffusional recycling of the first redox reversible species. Optionally the
10 potential on one working electrode can be set at a first diffusible species dependent, anodic potential and current flow is measured as the potential of the other working electrode is swept through a potential corresponding to the predetermined diffusible species dependent cathodic potential (or *vice versa*).

The cathodic and anodic potentials appropriate for each reversible
15 redox species can be readily determined by empirical measurement. The multiple redox reversible species used in performance of the method of this invention are selected to have redox potentials differing by at least 50 millivolts, more preferably at least 100 millivolts, more preferably at least 200 millivolts, from that of each other redox reversible species utilized in the method. The difference in redox potentials of
20 the redox reversible species being used allow each species to be detected without significant interference from the second or any other redox reversible species in the liquid sample. A steady state current flow is rapidly established at each of the working electrodes following application of the anodic and cathodic potentials. Current flow can be measured at either or both working electrodes, and it is proportionate to the
25 concentration of the recycling redox reversible species.

Second cathodic and anodic potentials are applied to the working electrodes wherein said second potentials correspond to those respective potentials necessary to establish current flow through the sample due to diffusional recycling of the second redox reversible species without significant interference from the first redox
30 reversible species, and the resulting steady state current flow is measured. This step is repeated for each redox reversible species utilized in the method. The measured current flows are then correlated to known concentrations of the respective diffusible

redox reversible species. Those concentrations are proportionate to the respective analyte concentrations.

The method steps can be conducted using a programmed bipotentiostat to control potentials on the electrode structure in contact with the sample. The
5 bipotentiostat can be included either in a desktop or hand-held meter further including means for reading values for steady state current, storing said values, and calculating analyte concentrations using a microprocessor programmed for making such calculations.

The relative amounts of the first and second redox reversible species
10 and the respective specific binding partners for the targeted analytes to be measured in the method can be determined empirically. They are dependent on the concentration ranges of the targeted analyte, and the binding stoichiometry of the specific binding partner, the binding constant, the analyte and the corresponding redox reversible species. The amounts of each reagent appropriate for each analyte being measured can
15 be determined by empirical methods.

The present osmium conjugates can also be used in a method for measuring two proteinaceous analytes in a liquid sample wherein the ligand analog component of the first redox reversible species is a peptide comprising an epitope of a first analyte and the ligand analog component of a second redox reversible species is a
20 peptide comprising an epitope of a second analyte. One specific binding partner utilized in the method is an antibody recognizing the epitope of the first analyte, and the other specific binding partner is an antibody recognizing the epitope of the second analyte. In another application of that method two independent measurements are performed on a single analyte in a liquid sample. In that embodiment the respective
25 ligand analog component of the first and second redox reversible species are different ligand analogs of the targeted analyte. Where the targeted analyte is a proteinaceous compound, the ligand analog component of the first redox reversible species is a peptide comprising a first epitope of the analyte, and the ligand analog of the second redox reversible species is a peptide comprising a second epitope of the analyte, and
30 the specific binding partners are first and second antibodies, each recognizing respective first and second analyte epitopes.

The present osmium conjugates can also be used in a device for detecting or quantifying one or more analytes in a liquid sample. The device comprises at least two redox reversible species, each capable of diffusion in said liquid sample at least in the presence of a respective predetermined analyte, said redox reversible species having respective redox potentials differing by at least 50 millivolts, an electrode structure for contact with the liquid sample in said chamber, said electrode structure including a reference electrode and working electrodes dimensioned to allow diffusional recycling of a diffusible redox reversible species in a liquid sample in contact with the electrode system when a predetermined redox-reversible-species-dependent cathodic potential is applied to one working electrode and a predetermined redox-reversible-species-dependent anodic potential is applied to a second working electrode, said diffusional recycling of said species being sufficient to sustain measurable current through each working electrode, and conductors communicating with the respective electrodes for applying said anodic potential and said cathodic potential and for carrying the current conducted by the electrodes.

The device can be constructed using procedures and techniques that have been previously described in the art for construction of biosensors employing electrometric detection techniques. Thus, for example, the device can include a chamber that has a receiving port, and the chamber is dimensioned so that it fills by capillary flow when the liquid sample is contacted with the sample receiving port. The electrode structure can be formed on a plate that defines a wall of the chamber so that the electrode structure will contact a liquid sample in the chamber. Thus, for example, the device can be constructed using the general procedures and designs described in U.S. Patent No. 5,141,868, the disclosure of which is expressly incorporated herein by reference. The features of the present invention can also be incorporated into other electrochemical biosensors or test strips, such as those disclosed in U.S. Patent Nos. 5,120,420; 5,437,999; 5,192,415; 5,264,103; and 5,575,895, the disclosures of which U.S. Patents are expressly incorporated herein by reference. The device can be constructed to include the predetermined amounts of the redox reversible species and the specific binding partners. For example, a mixture of such reagents can be coated onto a wall of the sample chamber in said device during device construction, so that

the liquid sample is contacted with the reagent mixture as it is delivered into the chamber for containing the sample. In one embodiment the device is constructed for quantifying a first analyte and a second analyte in liquid sample. The device comprises two redox reversible species, a first redox reversible species comprising a conjugate of
5 a ligand analog of the first analyte and a second redox reversible species comprising a conjugate of a ligand analog of the second analyte, and a specific binding partner for each analyte so that each of said analyte analog conjugates are capable of binding competitively with its respective analyte to a specific binding partner.

In another application of the present osmium conjugates, they are used
10 in conjunction with a device that further comprises a bipotentiostat in electrical communication with the conductors for applying a redox- reversible-species-dependent-cathodic potential to one working electrode and a redox-reversible-species-dependent-anodic potential to a second working electrode. The biopotentiostat can be programmed to apply a sequence of potentials to the respective working electrodes.
15 More particularly, the bipotentiostat can be programmed to apply first cathodic potential to a first working electrode and a first anodic potential to a second working electrode, said first anodic and cathodic potentials corresponding to those potentials necessary to establish current flow to the sample due to diffusional recycling of the first redox reversible species. The bipotentiostat is also programmed to apply a second
20 cathodic potential to said first working electrode and a second potential to the second anodic electrode, said second cathodic and anodic potentials corresponding to those potentials necessary to establish current flow through the sample due to diffusional recycling of the second redox reversible species. In an alternate embodiment the device includes first and second redox reversible species, and at least first and second
25 electrode structures for contact with the liquid sample in the chamber, each of said electrode structures comprising a microarray of working electrodes, and means for switching the bipotentiostat between the first and second electrode structures. In preferred device embodiments there is provided means for measuring current flow through the sample at each of the first and second potentials and preferably storing
30 values for said current flows in a register coupled to a microprocessor programmed to calculate analyte concentrations based on said values.

In still another embodiment the present conjugate is used in a kit for measuring the concentration of one or more analytes in liquid sample. The kit comprises

- at least two redox reversible species for contact with the liquid sample, each capable of diffusion in the liquid sample at least in the presence of a predetermined analyte, at least one of such species being a conjugate of a ligand analog of an analyte and a redox reversible label, said redox reversible species having respective redox potentials differing by at least 50 millivolts;
- a specific binding partner for each analyte;
- an electrode structure for contact with the liquid sample, said electrode structure including a reference electrode and working electrodes dimensioned to allow diffusional recycling of diffusible redox reversible species in the sample when a predetermined redox-reversible-species-dependent-cathodic potential is applied to one working electrode and a predetermined redox-reversible-species-dependent-anodic potential is applied to the second working electrode, said diffusional recycling of said species means sufficient to sustain a measurable current through the sample; and
- conductors communicating with the respective electrodes for applying said anodic potential and said cathodic potential and for carrying the current conducted by the electrodes.

- In one embodiment, the present redox reversible conjugate species are mixed with other electroactive species as a novel composition for contact with the liquid sample. In another embodiment each of the redox reversible species and the specific binding partner for each analyte is mixed as a novel composition for contact with the liquid sample. Preferably, the redox reversible label of at least one of the redox reversible species comprises an osmium complex of this invention.


Preparation of Os Mediator Labels

- The Os mediator *bis*(bipyridyl) imidazolyl chlorooxmium has been shown to be an excellent electron mediator for many oxide-reductase enzymes (U.S. Patent No. 5,589,326). It has fast mediation kinetics (about 500 times faster than ferricyanide with glucose oxidase) and a relatively low redox potential (+150 mV vs. Ag/AgCl). It has also very fast electron transfer rate at electrode surface. More



importantly, the organic ligands on Os mediator can be functionalized so that it can be covalently linked to other molecules without detrimental effects on redox properties of the Os center. These unique properties of Os mediator make it an ideal electrochemical indicator for sensors based on immunoaffinity.

5 Os mediators with these new ligands were synthesized using the same procedure used for Os free mediator. Their synthesis consists of two major process steps as outlined below. Details of these processing steps are described below.

The first process step involves the synthesis of Os intermediate, *cis*-bis(2,2'-bipyridyl) dichloroosmium(II), from commercially available osmium salt using
10 the following scheme. The intermediate product is isolated through recrystallization in an ice bath.





15



20

The second process step involves the reaction between Os intermediate and histamine or 4-imidazoleacetic acid (or a substituted bipyridine for preparation of the *tris*(bipyridyl) complexes) to produce Os mediators with the appropriate "handle". The desired product is then precipitated out from solution by addition of ammonium
25 tetrafluoroborate.



These Os mediators can also be easily converted to oxidized form, *i.e.* Os(III) using nitrosonium tetrafluoroborate. However, this is unnecessary since the Os revert back to reduced form anyway at alkaline conditions during conjugation reactions. And it does not require oxidized form of Os(III) for the detection on the
5 biosensor.

A. Simple Mixed Mediator Measurement

1. Interdigitated array microelectrodes (IDA) are produced through photolithographic means by art-recognized methods, (See WO 97/34140; EP
10 0299,780; D.G. Sanderson and L.B. Anderson, *Analytical Chemistry*, 57 (1985), 2388; Koichi Aoki et al., *J. Electroanalytical Chemistry*, 256 91988) 269; and D. Niwa et la., *J. Electroanalytical Chemistry*, 167 (1989) 291. Other means which are standard in lithographic processing may also be used to produce the desired patterns of a conductor on insulator substrate.

15 2. Reversible mediators are selected from those described herein and those described references (U.S. Patent Nos. 4,945,045 and 5,589,325, the disclosures of which are incorporated herein by reference). Preferably two different mediators are selected with potentials which differ by at least 100 mV, more preferably at least 200 mV. Examples of suitable mediators include the Os(bipy)₂ImCl of this
20 invention and in U.S. Patent No. 5,589,326, the disclosure of which is incorporated herein by reference, and ferrocene, described in U.S. Patent No. 4,945,045 and EP 0142301, the disclosures of which are incorporated herein by reference. Mixtures of these mediators are made in aqueous solution, for example phosphate-buffered saline (PBS). Concentrations between about 1 uM and 1000 uM may conveniently be
25 measured.

3. The IDA is connected to a bipotentiostat, an instrument capable of controlling the potential of two separate electrodes. Also provided is a reference electrode. This non-polarizable electrode serves as the reference for the two applied potentials and may also serve as the counter electrode. Any non-polarizable electrode
30 may be used, for example Ag/AgCl, such as may be obtained from ABI/Abtech. An auxiliary electrode can also be used for controlling current flow through the working electrodes. The mixtures are placed on the IDA electrode and the reference electrode

also contacted with the mixture, or the IDA along with the reference electrode may be dipped into the mixture.

4. To measure Mediator 1 ($\text{Os}(\text{bipy})_2\text{ImCl}$)

A cathodic potential is applied to one set of fingers of the IDA which is capable of
5 reducing mediator 1 (ca -50 mV vs. Ag/AgCl). An anodic potential is applied to the
other set of fingers of the IDA which is capable of oxidizing mediator 1 but not
mediator 2 (or any other mediators) (ca 250 mV vs Ag/AgCl). After a short time
(msec to sec), a steady state current will be measurable which is dependent only on the
concentration of mediator 1.

10 5. To measure Mediator 2 (Ferrocene)

A cathodic potential is applied to one set of fingers of the IDA which is capable of
reducing mediator 2 but not mediator 1 (ca 250 mV vs. Ag/AgCl). An anodic
potential is applied to the other set of fingers of the IDA which is capable of oxidizing
mediator 2 (ca 550 mV vs Ag/AgCl). After a short time (msec to sec), a steady state
15 current will be measurable which is dependent only on the concentration of mediator 2.

Specific Binding Assay with Mixed Mediator Measurement.

Specific Assay of HbA1c in a blood sample

20 1. IDA electrodes are provided as in Paragraph A above.
2. Conjugates of mediators 1 and 2 and haptens or specific binding
members are provided using art-recognized procedures for covalent coupling using
either a homo-functional or hetero-functional linker. Specifically, a synthetic peptide
corresponding to the N-terminal sequence of the β -chain of HbA1c is conjugated to the
25 osmium complex. Similarly, a synthetic peptide corresponding to the N-terminal
sequence of HbA0 is conjugated to a second mediator, for example ferrocene.

3. Antibodies for the analytes (HbA1c and HbA0) which react
specifically with the N-terminal peptides which have been incorporated into the
conjugate are provided by standard methods for producing polyclonal antibodies. In
30 this case, sheep were immunized with carrier proteins to which were conjugated the
synthetic peptide sequences for HbA1c and HbA0. Following the appropriate
immunization schedule, the sheep were bled, and the antibody isolated from the blood
via ion exchange chromatography, followed by immunosorbent purification on a
column of the same N-terminal peptide with a different linker.

4. Appropriate stoichiometry of the reaction was determined for the two reactions independently by methods standard for immunoassay development. A solution containing a fixed amount of labeled conjugate was mixed with a solution with varying amounts of antibody, and, following an appropriate incubation period, the amount of free conjugate remaining was measured on the IDA electrode using the procedure described above. The amount of antibody just sufficient to achieve maximum inhibition of the conjugate (ca > 80%) was selected.

5. Reagent solution 1 was made containing a mixture of the two conjugates in the appropriate concentrations. Reagent solution 2 was made containing a mixture of the two antibodies in the amounts determined above. A blood sample was diluted ca 20-fold in a solution of 25mM citric acid/0.5% Brij-35. Following a 30 second incubation to allow for lysis and denaturation of the hemoglobin, to 66 uL of this diluted sample was added 33 uL of 1 M phosphate buffer, to adjust the pH back to neutral. 30 uL of antibody solution 2 was added, and the mixture allowed to incubate 30 sec. Then 30 uL of conjugate solution 1 was added, and the mixture measured on the IDA electrode. The concentration of HbA1c in the sample is related to the current measured from Mediator 1, and the concentration of HbA0 is related to the current from Mediator 2. The %HbA1c in the sample is related to the ratio of the measured amounts of Mediator 1 and Mediator 2.

Application to HbA1c Assay

Hemoglobin A1c is a specific glycohemoglobin in which the glycosylation takes place at the N-terminal of hemoglobin b-chain. The antibody binds specifically to HbA1c has an epitope sequence of Gluc-Val-His-Leu-Thr. To facilitate conjugation to other molecules, a nonnative amino acid has been added to the sequence, *e.g.*, Cys, Lys, or Lys-MH, to produce A1c peptides including: 1) Gluc-Val-His-Leu-Thr-Lys-MH; 2) Gluc-Val-His-Leu-Thr-Lys; 2) Gluc-Val-His-Leu-Thr-Cys.



HbA1c assay requires measuring both A1c concentration and total hemoglobin concentration and reports the results as a ratio of these two measurements (%HbA1c). It is advantageous to assay both A1c and total hemoglobin using same principle because ratioing would minimize biases due to environmental effects. Thus antibody has been raised to bind specifically to hemoglobins with unglycosylated N-

terminus, *i.e.* with an epitope sequence of Val-His-Leu-Thr. Similarly, nonnative amino acid is added to the sequence to facilitate conjugation. The peptides used for total hemoglobin measurement is termed as A0 peptide. A0 peptides that have been used in the preparation of Os mediator-peptide conjugates include Val-His-Leu-Thr-
5 Cys and Val-His-Leu-Thr-Lys.

Conjugation Chemistry and Conjugates

There are many types of conjugation chemistry that can be employed to link Os mediator to a peptide. The following two conjugation chemistries employed
10 for the preparation of Os mediator-peptide conjugates have also been commonly used for preparing protein conjugates: 1) formation of amide bond by reactive ester with primary amine; 2) formation of thioether bond by maleimide with sulfhydryl group. Amide bond is preferred over thioether bond because amide bond is generally more stable. Based the preferred conjugation chemistry, the ligand on Os mediator can be
15 functionalized with either a primary amine group or a carboxylic acid group. The best location for these functional groups is believed to be the C-4 or C-5 positions on the imidazole ligand of Os mediator, however, functionalization through the non-Os-complexed imidazole ring nitrogen atom can also be carried out. Two different functionalized Os mediators were synthesized as described above.

20



25

Os mediator (a) was formed with histamine while Os mediator (b) was formed with imidazolacetic acid. However, it was found that the imino nitrogen of the


imidazole ring interferes with the activation of carboxylic acid group to reactive ester (*i.e.*, N-hydroxysuccinimide ester) using carbodiimide. Thus, use of carboxylic acid functionalized Os mediator in the synthesis of Os mediator-peptide conjugates gave much less favorable results.

5 The amine group on histamine ligand of Os mediator readily reacts with N-hydroxysuccinimide (NHS) ester to form amide bond. Two types of crosslinkers have been employed to link Os mediator to peptides, (a) heterofunctional crosslinker, having a NHS ester at one end and the other end has a maleimide or a sulfhydryl group; and (b) homofunctional crosslinker, *e.g.* both ends have NHS esters.

10 In the case of heterofunctional crosslinker, the crosslinker is first reacted with Os mediator with histamine ligand (Os histamine) at 1:1 molar ratio. One particular point needs to be noted here. Os mediator in oxidized form, *i.e.* Os(III), can instantly oxidize sulfhydryl group to form disulfide bond. It is important to keep Os center in the reduced form by addition of a small amount of reductant such as sodium
15 dithionite during the conjugation processes. The reaction progress can be monitored by analytical reverse-phase HPLC on a C18 column. Then the Os mediator-crosslinker adduct is isolated via preparative HPLC and the collected fraction is subsequently freeze-dried. Finally, the Os mediator-crosslinker adduct is reacted with the appropriate peptide to form Os mediator-peptide conjugate. Again, the product is
20 isolated by collecting appropriate fraction in preparative HPLC and the collected fraction is then freeze-dried.


Two different heterofunctional crosslinkers have been used for the synthesis of Os mediator-peptide conjugates. SMCC (succinimidyl 4-[N-maleimidomethyl]cyclohexane-1-carboxylate) is used for cystein-containing peptide,
25 while SATA (N-succinimidyl S-acetylthioacetate) is used for maleimide-containing peptide. Three Os mediator-peptide conjugates (two with A1c peptide and one with A0 peptide) have been made using heterofunctional crosslinkers and their structure are shown below: (a) Os-SMMC-A1c; (b) Os-SATA-A1c, and (c) Os-SATA-AO.

-27-




(a)

5



(b)

10




(c)

However, it has been found that these conjugates were not stable when they were stored as solutions. Analytical HPLC results indicated that these conjugates degrade. Mass spectroscopy confirmed that the instability is due to splitting of thioether bond present in these conjugates.

In order to avoid thioether bond in the conjugate, homofunctional crosslinker containing two NHS esters was used instead to prepare the conjugates.

The crosslinker used was DSG (disuccinimidyl glutarate). In order to prevent the formation of crosslinked Os mediator, *i.e.* Os-crosslinker-Os, a large excess of homofunctional crosslinker was used in the reaction with Os histamine at 4:1 molar ratio. Under this condition, only the desired product, *i.e.* Os-crosslinker, was formed.

- 5 The Os-DSG-A1c conjugate was similarly prepared using the procedure described earlier.



- 10 The preparation of analogous Os-DSG-A₀ conjugate requires an extra step since the unglycated N-terminal amine of HbA₀ peptide is also reactive toward NHS ester. In this case, the N-terminal amine of HbA₀ peptide is first protected by either a base-labile Fmoc¹ or an acid-labile Boc² group. After reacting with Os-DSG adduct to form Os-DSG-A₀ conjugate, the protecting group is then cleaved using
- 15 appropriate deprotection method (adding base for Fmoc or acid for Boc). The peptides prepared by solid-phase peptide synthesis already have N-terminal Fmoc protecting groups. The protecting groups are usually removed prior to cleavage of peptide from the resin beads, but they can also be left on if so desired. The HbA₀ peptide from Zymed has an intact Fmoc protecting group at N-terminal. Using this
- 20 strategy the Os-DSG-A₀ conjugate was successfully synthesized.

Many analytes cannot be assayed using enzyme-based sensors. They require the development of affinity biosensors or immunosensors which are based on the selective binding of antigens to antibodies. The key to the detection of this binding

¹Fmoc = 9-fluorenylmethyloxycarbonyl

²Boc = t-butyloxycarbonyl

event on electrochemical sensors is the inclusion of antigens labeled with redox labels. Bis(2,2'-bipyridyl) imidazolyl chloroosmium, a.k.a. Os mediator, possesses many properties that make it an excellent redox label for this purpose. In addition, a "handle" for linking it covalently to antigens can be added on without affecting its redox properties.

Several assay schemes can be used in affinity biosensors including, i) competitive binding assay (labeled antigen is competing for a limited number of binding sites); ii) sequential binding assays (labeled antigen is bound to excess binding sites); iii) heterogeneous assay (uses a separation step to separate bound and free labeled antigens); and iv) homogeneous assay (no separation step). The steps involved in a homogeneous sequential binding assay include binding the analyte to an antibody. The labeled antigen (analyte analog) binds to the remaining binding sites on the antibody. Finally the leftover free labeled antigen is detected at electrode surface. The resulting current will be a function of the amount of analyte present.

The detection of free labeled antigens can be achieved using either direct detection or amplified detection methods. Direct detection requires the use of advanced electrochemical techniques such as ac voltammetry, differential pulse voltammetry, square wave voltammetry or ac impedance in order to reach a sensitivity of 5 μ M or less. Amplified detection methods use dc amperometry with amplification through reaction with enzyme or chemical reductants or by using interdigital array (IDA) microelectrodes. The preferred detection method is amplified amperometry through cycling of free Os mediator label by using IDA microelectrodes. However, amplified amperometry using Gluc-DOR enzyme can also be used. Fast mediation kinetics of Os mediator is very desirable because the magnitude of amplification is dependent on its mediation kinetic constant with the enzyme.

General Analytical HPLC Method For Osmium Conjugates

All HPLC analysis were performed using a Beckman System Gold HPLC system consists of a 126 pump module and a 168 diod array detector module. Stationary phase is a Vydac analytical reverse-phase C18 analytical column. Other parameters are listed below.

-30-

Mobil Phase: A = 0.1%TFA³ in H₂O
 B = 0.1% TFA in CH₃CN

Flow rate: 1 mL/min

Gradient: 0- 5 min: 10% B
 5-45 min: 10% B -> 50% B at 1%/min
 45-50 min: 100% B

Detector: Channel A at 384 nm
 Channel B at 220 nm.

Synthesis of Bis(2,2'-bipyridyl) dichloroosmium

1. Charge a 1 L one-neck RB flask with 19.335 grams K₂OsCl₆ (0.04019 mole) and 13.295 grams 2,2'-dipyridyl (0.08512 mole). Add 400 mL DMF to dissolve all reactants.
2. Heat the solution to reflux and then maintain reflux for 1 hour. Then turn off the heat and let solution cool to 30-40°C at ambient.
3. Filter the reaction mixture using a medium grade glass-frit filter. Rinse the flask with additional 20 mL DMF and wash the filter.
4. Transfer the filtrate into a 3-L beaker. Charge another 2-L beaker with 22.799 grams of Na₂S₂O₄ and dissolve in 2 L deionized water. Add this solution to the beaker containing Os/DMF filtrate dropwise using a dropping funnel. Keep the solution stirring at all time.
5. Then cool the mixture in an ice bath for at least 3 hours. Add ice as necessary.
6. Filter the mixture "cold" using a ceramic filter with filter paper. Wash the content on the filter with 50 mL, water twice and 50 mL ether twice.
7. Dry the product under high vacuum at 50°C overnight (at least 15 hours). Weigh the product and transfer into a brown bottle. Store in a desiccator at room temperature.

Typical yield = 16 gram or 70%.

Product is analyzed by UV-Visible spectroscopy and elemental analysis.

UV-Vis: Peak λ (nm) ϵ (M⁻¹cm⁻¹)

³TFA = trifluoroacetic acid.

-31-

382	10,745
465	9,914
558	11,560
836	3,873

5

EA: C% H% N% Cl% Os% H₂O%

	Theoretical	41.89	2.81	9.77	12.36	33.17	0
10	Actual	40.74	2.92	9.87	11.91		0.41

Synthesis of Bis(2,2'-bipyridyl) histamine chloroosmium

- 15 1. Charge a 2L one-neck RB flask with 11.3959 gram Os(bpy)₂Cl₂ (0.0198 mole) and 4.9836 gram histamine (0.0448 mole). Add 500 mL ethanol to dissolve the reactants. Then add 250 mL deionized water.
- 20 2. Heat the solution to reflux and maintain reflux for 6 hours. Let solution cool to RT at ambient.
- 25 3. Remove all ethanol using rotary evaporation. Then transfer the solution into a 500 mL beaker. Dissolve 2.136 gram NH₄BF₄ in 20 mL water. Add dropwise to Os solution. Precipitate forms. Cool in an ice bath for 30 min. Filter the mixture using a ceramic filter with filter paper. Wash the content on filter with ~20 mL water twice.
- 30 4. Dry under high vacuum at 50°C overnight (at least 15 hour).
5. Weigh the product and transfer to a brown bottle. Store in a desiccator at room temperature.

Typical yield = 7.6 gram or 52%.

Product is analyzed by UV-visible spectroscopy and HPLC.

35

<u>UV-Vis:</u>	Peak λ (nm)	ε (M ⁻¹ cm ⁻¹)
	355	7,538
	424	7,334
	514	7,334
40	722	2,775

HPLC: Elution time = 18.0 min
Purity by HPLC range from 65-85%

45

Preparation of [Os(bpy)₂(histamine)Cl]-heterofunctional crosslinker adduct

-32-

1. Weigh 0.1167 g $[\text{Os}(\text{bpy})_2(\text{histamine})\text{Cl}]\text{BF}_4$ (0.162 mmol) and transfer to a 5 mL Reacti-Vial. Add 1.0 mL DMF to dissolve the reactant. Add 25 μL triethylamine.
- 5 2. Add 0.0508g SMCC(0.150mmol)or 0.0390g SATA (0.168mmol). Stir the reactants at RT for 2 hours. Inject a sample into HPLC to monitor reaction progress.
- 10 3. If reaction is complete, dilute the solution with 0.1% TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC and collect the product peak.
4. Freeze dry the collected fraction overnight.
- 15 5. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C .

Typical yield = 40 mg or 25%.

- 20 Product is analyzed by HPLC and ES/MS.

	<u>HPLC</u>	<u>ES/MS</u>
25 Os-SMCC:	Elution time = 32.1 min	$m^+/e = 434.2$
Os-SATA:	Elution time = 27.5 min	$m^+/e = 382.8$

30 **Preparation of $[\text{Os}(\text{bpy})_2(\text{histamine})\text{Cl}]\text{-homofunctional crosslinker adduct}$**

1. Weigh 0.2042 g DSG (0.626 mmol)) and transfer to a 5 mL Reacti-Vial. Add 0.75 mL DMF to dissolve the reactant.
- 35 2. Weigh 0.1023 g $[\text{Os}(\text{bpy})_2(\text{histamine})\text{Cl}]\text{BF}_4$ (0.142 mmol) and transfer to a separate 5 mL Reacti-Vial. Add 1.0 mL DMF to dissolve the reactant. Add 25 μL triethylamine. Then add Os/DMF solution dropwise to DSG/DW solution with constant stirring. After reacting for 2 hours at RT, inject a sample into HPLC to monitor reaction progress.
- 40 3. If reaction is complete, dilute the solution with 0.1% TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC and collect the product peak.
- 45 4. Freeze dry the collected fraction overnight.

5. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C.

Typical yield = 45 mg or 35%.

- 5 Product is analyzed by HPLC and ES/MS.

	<u>HPLC</u>	<u>ES/MS</u>
Os-DSG:	Elution time = 27.1 min	m ⁺ /e = 429.2 and 859.6

10

Preparation of Os-SATA-Alc Conjugate

- 15 1. Weigh 40.5 mg Os-SATA (0.0529 mmol) and transfer to a 5 mL Reacti-Vial with stir bar. Add 1.0 mL PBS (pH 7.5) to dissolve. Add 20 mg Na₂S₂O₄ in order to keep Os in reduced form.
- 20 2. Add 1.0 mL deacetylation buffer (PBS pH7.5 + 0.5 M hydroxylamine and 25 mM EDTA) to deprotect the sulfhydryl group. Inject a sample into analytical HPLC to determine whether deprotection is complete by appearance of a new peak at 25.8 min.
- 25 3. Add 45 mg HbAlc-MH peptide (0.0474 mmol) and let react at RT for 1 hour. Inject a sample into analytical HPLC to monitor reaction progress.
- 30 4. If reaction is complete, dilute the mixture with 0.1 %TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC to collect product peak.
5. Freeze dry the collected fraction overnight (at least 15 hour).
- 35 6. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C.

Typical yield = 12 mg

Product is analyzed by HPLC and ES/MS.

	<u>HPLC</u>	<u>ES/MS</u>
Os-SATA-Alc:	elution time = 27.6 min	m ⁺ /e = 559.1 and 838.5

45

Preparation of Os-SMCC-Alc Conjugate

-34-

1. Weigh 39.0 mg Os-SMCC (0.0452 mmol) and transfer to a 5 mL Reacti-Vial with stir bar. Add 1.0 mL PBS (pH 6.0) to dissolve.
- 5 2. Add 30.0 mg Hblc-Cys peptide (0.0450 mmol). Let reaction proceed at RT for 2 hours. Inject a sample into analytical HPLC to monitor reaction progress.
- 10 3. If reaction is complete, dilute the mixture with 0.1% TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC to collect product peak.
4. Freeze dry the collected fraction overnight (at least 15 hour).
- 15 5. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C.

Typical yield = 12 mg

Product is analyzed by HPLC and ES/MS.

20

HPLCES/MS

Os-SMCC-Alc: elution time = 27.6 min $m^+/e = 480.5$ and 534.4

25

Preparation of Os-SMCC-Ao Conjugate

- 30 1. Weigh 37.0 mg Os-SMCC (0.0426 mmol) and transfer to a 5 mL Reacti-Vial with stir bar. Add 1.0 mL PBS (pH=6.0) to dissolve.
- 35 2. Add 24.3 mg HbAo-Cys peptide (0.0425 mmol). Let reaction proceed at RT for 2 hours. Inject a sample into analytical HPLC to monitor reaction progress.
- 40 3. If reaction is complete, dilute the mixture with 0.1%TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC and collect the product peak.
4. Freeze dry the collected fraction overnight (at least 15 hour).
5. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C.

45 Typical yield = 15 mg

Product is analyzed by HPLC and ES/MS.

HPLCES/MS

Os-SMCC-A₀: elution time = 27.9 min m⁺/e = 360.7 and 720.5

5

Preparation of Os-DSG-Alc Conjugate

1. Weigh 32.0 mg Os-DSG (0.037 mmol) and transfer to a 5 mL Reacti-Vial with stir bar. Add 0.75 mL DMF to dissolve. Add 25 μ L triethylamine.
2. Add 26.5 mg HbAlc-Lys peptide (0.0349 mmol). Let reaction proceed at RT for 2 hours. Inject a sample into analytical BPLC to monitor reaction progress.
3. If reaction is complete, dilute the mixture with 0.1%TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC and collect the product peak.
4. Freeze dry the collected fraction overnight (at least 15 hour).
5. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C.

20

25

Typical yield = 16 mg

Product is analyzed by HPLC and ES/MS.

30

HPLCES/MS

Os-DSG-Alc: elution time = 23.5 min m⁺/e = 501.8 and 752.8

35 Preparation of Os-DSG-Ao Conjugate

1. Weigh 52.0 mg Os-DSG (0.0605 mmol) and transfer to a 5 mL Reacti-Vial with stir bar. Add 1.0 mL DMF to dissolve. Add 25 μ L triethylamine.
2. Add 49.1 mg Fmoc-HbA₀ peptide (0.0606 mmol). Let reaction proceed at RT for 2 hours. Inject a sample into analytical HPLC to monitor reaction progress by the appearance of peak at 40.3 min for Os-DSG-A₀(Fmoc).
3. If reaction is complete, inject additional 100 μ L triethylamine. After 1 hour, inject sample into analytical HPLC to determine whether all Fmoc protection group is removed by disappearance of the peak at 40.3 min.

40

45

4. If removal of Fmoc is complete, dilute the mixture with 0.1%TFA buffer to a final volume of 4.5 mL. Inject into preparative HPLC to collect product peak.
5. Freeze dry the collected fraction overnight (at least 15 hour).
6. Weigh the product and transfer to a brown bottle. Store in a desiccated bag at -20°C.

10 Typical yield = 16 mg

Product is analyzed by HPLC and ES/MS.

	<u>HPLC</u>	<u>ES/MS</u>
15 Os-DSG-A ₀ :	elution time = 23.2 min	m ⁺ /e = 447.4 and 670.3

20 **Synthesis of bis (4,4'-dimethyl-2,2'-bipyridyl) 4-methyl-4'-carboxylpropyl-2,2'-bipyridyl osmium [Os(dm-bpy)2(mcp-bpy)]Cl₂**

Potassium hexachloroosmium was reacted with 4,4'-dimethyl-2,2'-dipyridyl at 1:2 molar ratio by refluxing in DMF. The potassium chloride precipitate was filtered and the dimethyl-bipyridyl dichloroosmium complex was reduced from +3 oxidation state to +2 oxidation state using excess sodium dithionite. The product was recrystallized in DMF/water mixture at 0° and recovered by filtration.

4,4'-Dimethyl-2,2'-bipyridyl dichloroosmium was reacted with 4-methyl-4'-carboxylpropyl-2,2'-dipyridyl by refluxing in ethylene glycol. The solvent was removed by rotary evaporation. The product was dissolved in DMF and precipitate in ethyl ether. The product was dried in a vacuum oven overnight.

30 Analyticals: Product and intermediate product were analyzed by HPLC and mass spectroscopy for purity and identity of the compound.

Os(dm-bpy)2Cl₂: Theoretical MW = 629.6, MS showed 8 isotope peaks with most abundant peak at 630. HPLC elution time at 29.94 min with a purity of 90%+

35 Os(dm-bpy)2(mcp-bpy): MS confirmed the MW at 814.5 and HPLC showed a purity greater than 85%.

Synthesis of biotin-Os complex conjugate Biotin-Os(dm-bpy)₂(mcp-bpy)]Cl₂

The carboxyl group was activated by reacting the above Os complex with dicyclohexylcarbodiimide in the presence of N-hydroxysuccinimide. The active ester Os complex was isolated using preparative HPLC method and then reacted with
 5 amine-containing biotin to form the final conjugate.

Experiment to independently measure the concentration of two electroactive conjugate species.

10 Os(bipy)HisCl-DSG-HbA1c was prepared as described above.

Ferrocene-AMCHA-DADOO-biotin was prepared from ferrocene monocarboxylic acid, the crosslinker aminomethylcyclohexylic acid, the chain extender 1,8-diamino-3,6-DiOxoOctane and biotin as described elsewhere.

Mixtures of the two conjugates were prepared to evaluate the ability of
 15 the method of the invention to independently measure the concentration of the conjugates, and make corrections for variations in reagent amounts, electrode response, and environmental conditions.

Part 1: Simple Mixed conjugate response

20 The following matrix of solutions was prepared in 10 mM phosphate buffer with 150 mM NaCl and 0.5 % Brij-35, a non-ionic surfactant.

25

Os-DSG-HbA1c	Ferrocene-Biotin	Os-DSG-HbA1c	Ferrocene-Biotin
uM/l	uM/l	uM/l	uM/l
0	0	0	12.5
6.25	0	6.25	12.5
12.5	0	12.5	12.5
25	0	25	12.5

30

35

Os-DSG-HbA1c	Ferrocene-Biotin	Os-DSG-HbA1c	Ferrocene-Biotin
uM/l	uM/l	uM/l	uM/l
0	25	0	50
6.25	25	6.25	50

12.5	25	12.5	50
25	25	25	50

5 An Interdigitated array (IDA) microelectrode was fabricated according to the procedures described. In addition to the IDA, the chip had a silver/silver chloride electrode on the surface to function as the reference electrode and counter electrode. This electrode was produced with the same lithographic process, and then electroplated with silver and silver chloride according to standard techniques. The IDA
10 was connected to a bipotentiostat capable of controlling the potential relative to the reference and measuring the current at each of the electrodes of the IDA. Aliquots of the solutions were placed onto the surface of the chip, such that the IDA and the reference electrodes were covered.

Measurements were made by first applying -100 mV (vs. Ag/AgCl) to
15 one electrode of the IDA, and 200 mV to the other electrode for a period of 30 seconds. At this time, current was measured at each electrode. Then 200 mV was applied to one electrode, and 550 mV to the other. After 30 seconds, current was again measured. See Figure 9 for a summary of the results, which clearly demonstrate that the concentrations of the two mediators can be independently measured with this
20 method.

Part 2. Concentration Co-variance of Dual Mediators on IDA electrodes.

In this experiment, it was demonstrated that by making a mixture of known concentrations of two different mediators, and measuring different dilutions of
25 that mixture by the method of the invention, the ratio of the concentrations of the mediators remains constant. (Internal standard application).

The same two mediator conjugates were used as in Part 1. (Os-DSG-Alc and Fc-Bi)

From a solution containing 40 μ M of each conjugate, solutions
30 containing 27 μ M, 32 μ M, and 36 μ M of each conjugate were prepared in the same buffer (PBS/Brij).

The solutions were measured as in the previous example. Each solution was measured on 5 different IDA electrodes. The results are presented as the means,

standard deviations, and coefficient of variation for each solution separately, and for all solutions pooled over all electrodes.

5	Individual concentrations		Os-DSG-A1c	Fc-Bi	Os/FC
	28 uM	Mean	156	85	1.84
		S.D.	17.5	13.2	0.08
		% C.V.	11.2	15.5	4.4
10	32 uM	Mean	165	88	1.88
		S.D.	11	7.2	0.04
		% C.V.	6.7	8.2	2.0
	36 uM	Mean	189	102	1.85
		S.D.	12.5	6.9	0.04
		% C.V.	6.6	6.7	2.3
15	40 uM	Mean	208	110	1.90
		S.D.	9.5	5.5	0.06
		% C.V.	4.6	5.0	3.4
	Pooled Concentrations	Mean	182	97	1.87
		S.D.	24.4	13.2	0.06
20		% C.V.	13.4	13.5	3.4

This example clearly demonstrates that the internal standard effect of measuring two conjugates or mediators and calculating the ratio gives significantly improved precision of measurement, not only within each solution (compensation for variation between electrodes) but over all solutions (compensation for variation in sample dilution or amount).

Part 3. Temperature compensation

It was desired to show the effectiveness of the method in compensating for environmental influences such as Temperature variation on the accuracy or the measurement.

The same two conjugates were prepared in solution at 40 uM as before. They were measured as before on IDA electrodes, either at room temperature or warmed to 35-40 C on a heated metal plate prior to the measurement. The solutions were also warmed to 37 C prior to application to the electrodes.

	Room Temperature (23C)	Warmed (35-40) C	Ratio of response Warm / RT
Os-DSG-A1c	261	387	1.45
Fc-Bi	179	268	1.5
Ratio Os/Fc	1.46	1.44	0.99

5

As demonstrated by the results, the measured values increase by almost 50 % in the case of the warmed samples, which would lead to a large measurement error. However the use of the internal standard and ratio calculation effectively eliminates the temperature dependence of the result.

10

Immunoassay Detection of HbA1c with Osmium Mediator Conjugates.

The goal of all diabetic therapy is to maintain a near normal level of glucose in the blood. Home blood glucose kits are available to monitor the current glucose values and are valuable for diabetics to adjust day to day insulin doses and eating habits. Unfortunately, since the tests only measures a point in time result, it does not tell them the overall effectiveness of their actions in maintaining glycemic control. Measurement of glycosylated hemoglobin is now becoming widely accepted as an index of mean blood glucose concentrations over the preceding 6-8 weeks and thus provides a measure of the effectiveness of a diabetic's total therapy during periods of stable control. Since monitoring a diabetic's glycated hemoglobin can lead to improved glycemic control, the ADA recommends routine measurements of four times a year up to once a month for less stable type I diabetics.

Several technologies are available for the measurement of glycated hemoglobin. They include immunoassays for HbA1c (TinaQuant, BMC; DCA2000, Ames; and Unimate, Roche), ion exchange (Variant, BioRad; Eagle Diagnostics), and affinity chromatography (ColumnMate, Helena; GlyHb, Pierce).

One objective of this project is to develop a simple to use disposable strip for electrochemical detection of HbA1c for use in both physician offices and the home.

30

The most significant parameter for assessing patient condition is ratio of HbA1c to HbA₀, and thus the measurement of both glycated (HbA1c) and nonglycated (HbA0) values is required to calculate the ratio. This requires two separate measurements. It is preferable to use the same technology to measure both the

glycated and nonglycated fractions, thus removing some sample and environmental interferences. Measurement of HbAlc via electrochemical immunoassay is described below. Electrochemical HbA₀ immunoassay measurements are carried out using the same methods as that for HbAlc. The concentrations of HbA₀ are significantly higher.

- 5 One alternative to A₀ measurements using immunoaffinity would be to measure total hemoglobin directly using biamperometry or differential pulse voltammetry. This can be easily accomplished since hemoglobin is readily oxidized by $[Os(bpy)_2(im)Cl]^{2+}Cl_2$.

The N-terminal valine of the β -chain of hemoglobin A is the site of glycosylation in HbAlc, and serves as a recognition site for the antibody. In whole
10 blood the N-terminal valine is not accessible for the antibody to bind. Access is gained by lysing the red cells to release the hemoglobin followed by a conformation change (denaturing or unraveling) to adequately expose the HbAlc epitope. Dilution of the sample may occur as part of the lysing/denaturing process or may be required post denaturing to prepare the sample for the antibody (adjust pH, other) or bring the
15 sample into a range suitable for electrochemical immunoassay. In one embodiment, a fixed amount of antibody is incubated with the prepared sample and it binds to the HbAlc epitopes of the sample. The free antibody and the antibody bound sample is then combined with the osmium peptide conjugate (Alc or A₀) to allow the remaining unbound antibody to bind to the mediator label. When the mediator is bound to the
20 antibody (a macromolecule), it can not freely diffuse to interact with the electrode and thus currents generated are significantly reduced. The remaining unbound mediator label is therefore proportional to the concentration of HbAlc in the sample. The unbound mediator can be measured electrochemically either through an enzyme amplification method or directly using an interdigitated array electrode with
25 bipotentiostatic control.

An electrochemical HbAlc immunoassay response was demonstrated on gold electrodes using enzyme amplification in a biamperometric mode. The experimental conditions were not thoroughly optimized and the assay components were premixed in microcentrifuge tubes and then pipette onto the gold 4.95 mm² E-
30 cells. The experimental conditions for this assay are shown in Table 1. Table 2 shows the dose response data for the low, medium, and high HbAlc samples. Figure 4 shows the dose response with two additional points LL and HH which represent a diluted low

and a concentrated high sample. The higher currents for LL may possibly be explained by a reduction in blood proteins (due to dilution) leading to a reduction in electrode fouling.

5

Table 1. Dose Response Conditions

	Sample Preparation (denaturing)	Reagents	Electrochemical Measurement
10	1. 20 μ L Lysed blood for L, M, H samples (previously frozen whole blood)	1. 50 μ L 12.5 μ M PAB IS<Alc>in PBST	1. VXi Waveform E-450 mV for 60s
15	2. 18 μ L of L (LL) and 22 μ L of H for (HH)	2. 50 μ L denatured Blood	2. Insert Gold Electrodes WE = 1.5 x 3.3 mm
	3. 480 μ L 1.5 M LiSCN with 0.1% Tween	3. 20 μ L 40 μ M Os-DSG-Alc in DI H ₂ O	3. Apply 20 μ L Reagent
20	4. Mix and allow to denature for 10 minutes (Vortex)	4. 20 μ L 1 M Glucose in DI H ₂ O	4. Start biamprometric measurement mode
		5. 20 μ L Glucdor/PQQ (1.7mg/ml/0.17mg/ml)	5. Extract 60 sec data

Table 2 Blood Dose Response Data

Table 2: Data Response Data											
	Current @ 60 seconds (nA)				Separate denaturant step			All Data			
	g/L	#1	#2	#3	#4	Mean	SD	CV	Mean	SD	CV
5	6.49	76.28	78.34	84.79	78.47	79.47	3.69	4.64	81.54	4.67	5.73
	6.49	80.30	82.97	86.82	83.48	83.48	2.69	3.22			
	6.49	83.25	85.49	90.05	86.67	86.67	2.94	3.40			
	6.49	76.66	71.72	78.24	76.56	76.55	1.44	1.88			
10	12.34	95.61	95.05	93.06	94.02	94.44	1.13	1.20	98.31	4.19	4.26
	12.34	98.59	106.29	103.81	102.08	102.08	3.60	3.53			
	12.34	95.80	103.14	99.90	99.32	99.32	3.06	3.08			
	12.34	93.16	96.2	103.10	97.41	97.41	4.16	4.27			
15	18.20	100.06	110.83	119.08	112.52	112.52	9.29	8.26	113.48	5.08	4.48
	18.20	115.43	116.93	116.77	116.26	116.26	0.71	0.61			
	18.20	116.12	109.72	112.07	112.21	112.21	2.78	2.48			
	18.20	118.13	112.16	109.8	112.93	112.93	3.61	3.20			

	Calculated HbA1c (g/dL)				Separate denaturant step			All Data			
	g/L	#1	#2	#3	#4	Mean	SD	CV	Mean	SD	CV
20	6.49	4.46	5.22	7.58	5.26	5.63	1.35	24.00	6.39	1.71	26.80
	6.49	5.94	6.91	8.33	7.23	7.10	0.99	13.87			
	6.49	7.02	7.84	9.51	8.72	8.27	1.08	13.05			
	6.49	4.60	3.89	5.18	4.56	4.56	0.53	11.58			
25	12.34	11.55	11.34	10.61	10.97	11.12	0.41	3.72	12.54	1.54	12.25
	12.34	12.64	15.46	14.56	13.03	13.92	1.32	9.48			
	12.34	11.62	14.31	13.12	12.58	12.91	1.12	8.70			
	12.34	10.65	11.77	14.29	12.12	12.21	1.53	12.50			
30	18.20	13.18	17.13	20.15	20.54	17.75	3.41	19.19	18.10	1.86	10.29
	18.20	18.82	19.37	19.31	19.00	19.12	0.26	1.36			
	18.20	19.07	16.72	17.58	17.16	17.63	1.02	5.78			
	18.20	19.81	17.62	16.75	17.42	147.90	1.32	7.40			

n=16, 4 denaturation for each HbA1c sample x 4 replicates each, Y=2.82x+62.54, See file 4435051 A.xls

Blood lysis is necessary to release the hemoglobin followed by denaturing to expose the HbAlc epitope. Lysis can easily be accomplished via surfactants, osmotic effects of dilution with water, and directly by many denaturants. Blood lysed through a freeze/thaw cycle was shown not to significantly interfere with the biamperometric measurement ("open rate" with and without lysed blood was almost identical).
5 Conversely, denaturing the lysed blood with a variety of known denaturants to expose the HbAlc epitope has shown significant suppression of the electrochemical response, inhibiting measurement of an HbAlc dose response. Only LiSCN and citric acid from the list of evaluated denaturants shown in Table 3 was able to expose the Alc epitope
10 and minimize protein fouling enough to measure an HbAlc dose response.

Denaturing the sample for antibody recognition without severely fouling the electrode surface is important for successful development of an HbAlc immunoassay. Although LiSCN has been used almost exclusively to show feasibility, it has many limitations that would hinder its use in the disposable. Citric acid, a solid at RT may
15 offer benefits as a denaturant if it could be dried onto a strip followed by a diluent to adjust the pH to neutral. Acid or base blood denaturing followed by a final pH adjustment with a buffered diluent is an area worth further evaluation. One problem that was initially encountered was precipitation in adjusting the pH back to neutral, which can be overcome by using a different buffer or with the addition of surfactants.

Table 3. Blood Denaturants

	Method	Comments
	KSCN	<ul style="list-style-type: none"> Initial work did not show a dose response with KSCN denatured blood. KSCN has a larger negative effect on the electrochemical response than LiSCN. Literature shows that LiSCN is more effective than KSCN (concentrated efforts on LiSCN).
5	DCA2000 Buffer	<ul style="list-style-type: none"> Denaturing of blood not evident with the higher blood concentrations required for this assay. Higher concentrations of LiSCN are shown below.
	LiSCN	<ul style="list-style-type: none"> Method used by Ames DCA2000 HbA1c immunoassay.
10	Citric, Sulfuric, Hydrochloric, & Perchloric Acid	<ul style="list-style-type: none"> Blood HbA1c dose response (high/low) was seen with citric acid and was comparable to the response with LiSCN Evidence of blood denaturing was seen by all: "solution turned brown." Citric acid is preferred. Adjustment of pH to neutral after denaturing also saw problems of precipitation. Enzyme mediated responses with Gluc-Dor at pH 5.7 reduces response 50% compared to pH 7-8. Citric acid blood denaturing method is shown in Figure 5.
	Pepsin/Citric Acid	<ul style="list-style-type: none"> Roche HbA1c immunoassay uses pepsin/citric acid to hemolyze and proteolytically degrade hemoglobin to glycoproteins accessible by the antibody. Denaturation was apparent by the color change to a brownish red solution. Hemoglobin A1c dose response (high/low) was obtained comparable to LiSCN and citric acid denaturants. The procedure was identical to that of citric acid used above with the exception of pepsin added to the acid. Results were identical to that of citric acid.
15	TTAB (Tetra decyltrimethyl ammonium bromide)	<ul style="list-style-type: none"> Method of denaturing used in the TinaQuant HbA1c turbidimetric immunoassay. Evidence of denaturing: "solution turned green" TTAB concentrations 0.0125-0.2% severely suppressed enzyme mediated (Glucdor/PQQ/Glucose) biampereometric measurements. Open rates were 16-50 nA compared to 140 nA without TTAB.
	NaOH	<ul style="list-style-type: none"> Evidence of denaturing: "solution turned brown." NaOH does not adversely effect the enzyme mediated electrochemistry. Even at high pH the open rates do not change, although pH adjustment will probably be required to bring it within an optimal range for the antibody. NaOH denatured blood suppresses the open rates probably due to protein fouling. Lowering the pH to neutral tends to cause some precipitation.

Table 4. Effective Blood Denaturing Procedure for 2% Blood

LiSCN (One Step)	LiSCN (Two Step)	Citric Acid (2 Step)
40 μ L 6 M LiSCN 20 μ L 5% Tween in PBS 20 μ L Blood Mix (vortex) and allow to denature for 10 minutes. Dilute with 920 μ L DI H ₂ O.	960 μ L 1.5 M LiSCN 20 μ L 5% Tween in PBS 20 μ L Blood Mix (vortex) and allow to denature for 10 minutes.	200 μ L 0.2 M Citric Acid 20 μ L 5% Tween in PBS 20 μ L Blood Mix (vortex) and allow to denature for 10 minutes. Dilute with 760 μ L 8X PBS (0.1% Tween)
Denaturing time was not optimized. Limited data supports longer times for better precision using this method.	Denaturing time was not optimized. Data indicates shorter times may be adequate.	No optimization studies were performed.

PBS = 10 mM Phosphate Buffer, 2.7 mM KCl, 137 mM NaCl pH=7.4

Increased level of surfactant (5% Tween) reduces or eliminates precipitate.

Electrode fouling caused by denatured blood proteins adsorbing to the electrode surface can impede electron transfer and thus decrease electrode sensitivity. Electrode fouling or passivation occurs more or less immediately following sample contact with the surface thus minimizing the severity of denaturing in the sample should be the first approach. Surface conditions that are hydrophobic will favor adhesion of the proteins and thus fouling may be minimized with electrode surfaces of higher surface energies. This explains why gold electrodes shows less fouling with denatured blood than palladium. Reduction of protein fouling may be achieved by changing or protecting the electrode surface. Modifications that make the surface more hydrophilic should reduce the amount of fouling and can be accomplished by argon or oxygen plasma treatment or corona treatment. Selective coatings that could block the proteins from reaching the electrode surface can usually partially circumvent the problem have been used in the field to reduce fouling. Unfortunately, dramatic decreases in responses greater than seen with the denatured blood proteins are normally noted with their use. Hydrophilic coatings such as PEO were also evaluated and showed some improvement, but have similar problems of decreased magnitude and precision caused by forcing reagents to diffuse through the polymers. Reagents dispensed and dried over the electrodes may help reduce the magnitude of protein fouling with less negative effects.

Mediator concentration dose response, inhibition with antibody and reversal with a BSA-Alc polyhaptan were evaluated and summarized in Table 5. The Os-DSG-

Alc is stable in a lyophilized form and when frozen in solution at -20°C (40 and $80\text{ }\mu\text{M}$).

Table 5. Osmium Mediator Labels

5	Mediator Label	Concentration Response	Inhibition with Antibody	Reversal	Comments
	Os-SMCC-Alc	Linear	PAB IS ($\leq 92\%$) PAB DE ($\leq 97\%$) MAB ($\leq 50\%$)	Yes with BSA-Alc polyhaptan	% = inhibition values ranged from 16% to 97% depending on age of Os-SMCC stock solution. Degrades in solution.
10	Os-SATA-Alc	Linear	PAB IS ($\leq 44\%$)	Yes with BSA-Alc polyhaptan	Stability similar to SMCC.
	Os-DSG-Alc	Linear	PAB IS ($\leq 91\%$) PAB DE ($\leq 87\%$) MAB ($\leq 78\%$)	Yes with BSA-Alc polyhaptan	More stable than conjugate made with SATA and SMCC crosslinker but still degrades in solution.
	Os-SATA-A ₀	Linear	Yes with Sheep B<HbA ₀ > ($\leq 84\%$). No with Zymed rabbit antibodies	No with A ₀ HB-peptide#1	A ₀ conjugate was found to be unstable in solution. Conjugate was not lyophilized.

- 15 Polyclonal DE (ion-exchange) purified sheep antibody is used in the TinaQuant HbAlc assay. IS (immunosorbent) antibody is prepared using standard IS purification methodology. Samples of a monoclonal antibody were also obtained for evaluation. Inhibition curves were performed in solution with all mixing occurring in microcentrifuge tubes. Assays were measured by applying $20\mu\text{L}$ onto 6 mm^2
- 20 palladium electrodes with the conditions shown in Table 5. Inhibition curves with the three hemoglobin A1c antibodies (PAB IS, PAB DE, and MAB) were generated by fixing OS-DSG-A1c at $5\mu\text{M}$ and varying the antibody concentration. Both PAB IS and MAB showed the expected stoichiometric relationship for inhibition with the osmium peptide conjugate indicating efficient and fast binding of the antibody to the
- 25 A1c peptide. The polyclonal IS and monoclonal both showed steep inhibition curves with maximum change being reached close to $5\mu\text{M}$. Additional antibody above $5\mu\text{M}$ showed little effect on increasing the inhibition. The less purified PAB DE antibody had a much smaller slope and as expected required more than 3 times the amount to get close to maximum inhibition. Figure 6 shows the inhibition curves for each of the


HbAlc antibodies tested. From the inhibition curves we were able to select reasonable concentrations of antibody for maximum reversal with Alc samples.

Inhibition curves were also performed for the Os-SMCC-Alc (Max = 97%) and OS-SATA-Alc (Max = 44%) mediator labels. Stability of the mediator labels were
5 also evaluated by monitoring % inhibition values over time. All of the mediator labels showed some degradation when stored in dilute solutions (40 μ M) at RT. Samples frozen at -20°C appear to be stable.

For demonstrating inhibition reversal, antibody concentrations of 4 μ M for both PAB IS and MAB and 15 μ M for PAB DE were chosen from the inhibition
10 curves shown above. Reversal curves were then generated using a series of dilutions of BAS-Alc polyhaptan with a ~1:1 Alc:BSA. The BSA-Alc acts as our sample and binds to the antibody. Figure 10 shows the reversal curves for the three antibodies.

While these feasibility studies for a HbAlc immunoassay used an enzyme mediated amplification method. (Glucdor/PQQ/glucose was used to regenerate
15 reduced mediator after oxidation at the electrode surface providing a higher diffusion controlled current is given by the cottrell equation), they are considered to be indicative of results attainable with the use of IDA electrodes with bipotentiostatic control in accordance with this invention.

1. An electrochemically detectable compound of the formula



10 wherein

R and R₁ are the same or different and are 2,2'-bipyridyl, 4,4'-disubstituted-2,2'-bipyridyl, 5,5'-disubstituted-2,2'-bipyridyl, 1,10-phenanthrolyl, 4,7-disubstituted-1,10-phenanthrolyl, or 5,6-disubstituted-1,10-phenanthrolyl, wherein each substituent is a methyl, ethyl, or phenyl group,

15 R and R₁ are coordinated to Os through their nitrogen atoms;

q is 1 or 0;

R₇ is B-(L)_k-Q(CH₂)_i-;

R₂ is hydrogen, methyl, or ethyl when q is 1, and R₂ is B-(L)_k-Q(CH₂)_i-

when q is 0;

20 wherein in the group B-(L)_k-Q(CH₂)_i-

Q is O, S, or NR₄ wherein R₄ is hydrogen, methyl or ethyl;

-L- is a divalent linker;

k is 1 or 0;

i is 1, 2, 3, 4, 5 or 6; and

25 B is hydrogen or a group comprising a ligand capable of binding to a specific binding partner;

Z is chloro or bromo;

m is +1 or +2;

X is chloride, bromide, iodide, fluoride, tetrafluoroborate, perchlorate,

30 nitrate, sulfate, carbonate, or sulfite;

Y is chloride, bromide, iodide, fluoride, tetrafluoroborate, perchlorate

or nitrate; and

n is 1 or zero,

provided that when X is sulfate, carbonate, or sulfite, n is zero,

and when m is 1, n is zero and X is not sulfate, carbonate or sulfite.

2. The detectable compound of claim 1 wherein R and R₁ are the same or
5 different and are 2,2'-bipyridyl, 4,4'-disubstituted-2,2'-bipyridyl, 5,5'-disubstituted -2,2'-
bipyridyl.

3. The detectable compound of claim 2 wherein R₂ is hydrogen or methyl.

4. The detectable compound of claim 3 wherein Z is chloro.

5. The detectable compound of claim 4 wherein
10 X is chloride, bromide, or tetrafluoroborate,
Y is chloride, bromide, or tetrafluoroborate, and
when m is 2, n is 1, and when m is 1, n is zero.

6. The detectable compound of claim 1, wherein R and R₁ are each 2,2'-
bipyridyl.

7. The detectable compound of claim 1 wherein
15 R and R₁ are the same or different and are 2,2'-bipyridyl, 4,4'-dimethyl-
2,2'-bipyridyl, 4,4'-diphenyl-2,2'-bipyridyl, 1,10-phenanthroline, 4,7-dimethyl-1,10-
phenanthroline, 4,7-diphenyl-1,10-phenanthroline, or 5,6-dimethyl-1,10-
phenanthroline; and

20 R and R₁ are each coordinated to Os through their nitrogen atoms.

8. The detectable compound of claim 1, 2, 3, 4, 5, 6, or 7 wherein q is 1,
the group R₇ is bonded to C-4, and -L- is the divalent residue of a heterofunctional
crosslinker of the formula S-L¹-T wherein L¹ is a divalent linker and S and T are
different electrophilic groups capable of reacting with a nucleophilic group to form a
25 covalent bond.

9. The detectable compound of claim 1, 2, 3, 4, 5, 6, or 7 wherein q is 1,
the group R₇ is bonded to C-4, and -L- is the divalent residue of a homofunctional
crosslinker of the formula S-L¹-T wherein L¹ is a divalent linker and S and T are the
same electrophilic groups capable of reacting with a nucleophilic group to form a
30 covalent bond.

10. The detectable compound of claim 8 wherein B comprises an epitope
recognizable by an antibody capable of specific binding to a biologically significant

compound so that the antibody is capable of specific binding to said compound competitively with the biologically significant compound.

11. The detectable compound of claim 9 wherein B comprises an epitope recognizable by an antibody capable of specific binding to a biologically significant
5 compound so that the antibody is capable of specific binding to said compound competitively with the biologically significant compound.

12. The detectable compound of claim 11 wherein B comprises a glycosylated peptide of the formula

Gluc-Val-His-Leu-Thr-

10

and the compound binds competitively with glycosylated hemoglobin for an antibody whose epitope includes the glycosylated N-terminus of the β -chain of HbA_{1c}.

13. The detectable compound of claim 11 wherein B comprises a peptide of the formula

15

Val-His-Leu-Thr-

14. A method for measuring the concentration of an analyte in a sample of a liquid, said method comprising

- 20 contacting said sample with predetermined amounts of (1) a specific binding partner for said analyte, and (2) a detectable compound of claim 1 wherein B comprises a ligand selected so that the detectable compound binds competitively with the analyte to said binding agent, and

determining electrochemically the concentration of the detectable
25 compound of claim 1 not bound to the specific binding agent.

15. The method of claim 14 wherein the liquid sample is diluted blood, the analyte is glycosylated hemoglobin and the group B in the compound of claim 1 comprises a glycosylated peptide of the formula

30

Gluc-Val-His-Leu-Thr-

16. The method of claim 15 further including the step of determining the concentration of total hemoglobin and unglycosylated hemoglobin in the liquid sample, and calculating the ratio of the concentration of glycosylated hemoglobin to unglycosylated hemoglobin.

- 5 17. The method of claim 15 wherein the concentration of the unbound compound of claim 1 is determined by dc voltammetry using an interdigitated microarray electrode.

1/13




FIG. 1




FIG. 2





FIG. 3

3/13



n=16 for each HbA1c SAMPLE. (4 SAMPLE DENATURATIONS X 4 STRIP REPLICATES)

FIG. 4A





FIG. 5

5/13



6/13




FIG. 7

7/13




FIG. 8

8/13




FIG. 9

9/13





FIG. 10

10/13

 $C_{49}H_{56}N_{10}O_3S_0s$

F.W. 1055.30

FIG 11

11/13





FIG. 12

 $C_{67}H_{89}N_{15}O_{15}ClS$

M.W. 1602.24

FIG. 13




FIG. 14

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07F15/00 G01N33/58 G01N33/72 C07K5/10 C07K14/805

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07F G01N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 589 326 A (DENG ZHI D ET AL) 31 December 1996 (1996-12-31) cited in the application	1-7
A	the whole document	8-13
A	--- US 4 945 045 A (FORREST GORDON C ET AL) 31 July 1990 (1990-07-31) cited in the application example 1	1,8, 10-17
A	--- WO 93 25907 A (CENTRE NAT RECH SCIENT ;DEGRAND CHANTAL (FR); BLANKESPOOR RONALD () 23 December 1993 (1993-12-23) the whole document	1,8, 10-17
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 September 1999

Date of mailing of the international search report

22/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hart-Davis, J

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 14066 A (UNIV MANCHESTER ;TRELOAR PAUL HOWARD (GB); KANE JOHN WILLIAM (GB);) 23 June 1994 (1994-06-23) page 3, paragraph 4 -page 4, paragraph 2; claims 1-6 -----	1,8, 10-17
A	DE 43 44 646 A (SCHELLER FRIEDER PROF DR) 29 June 1995 (1995-06-29) example 1 -----	1,8, 10-17
A	WO 97 01097 A (SEREX INC) 9 January 1997 (1997-01-09) example 3 -----	1,8, 10-17
A	A J NIELSON, W P GRIFFITH: "Reactions of Osmium Tetroxide with Protein Side Chains and Unsaturated Lipids" JOURNAL OF THE CHEMICAL SOCIETY, DALTON TRANSACTIONS, 1979, pages 1084-1088, XP002114931 the whole document -----	1-17

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
US 5589326	A	31-12-1996	US	5846702 A	08-12-1998
US 4945045	A	31-07-1990	AT	100203 T	15-01-1994
			AU	597562 B	07-06-1990
			AU	4241185 A	09-01-1986
			CA	1268520 A	01-05-1990
			DE	3587720 D	24-02-1994
			DE	3587720 T	01-06-1994
			EP	0167248 A	08-01-1986
WO 9325907	A	23-12-1993	FR	2692357 A	17-12-1993
WO 9414066	A	23-06-1994	AT	157773 T	15-09-1997
			AU	693772 B	09-07-1998
			AU	5654794 A	04-07-1994
			CA	2150907 A	23-06-1994
			DE	69313664 D	09-10-1997
			DE	69313664 T	12-02-1998
			DK	672252 T	09-03-1998
			EP	0672252 A	20-09-1995
			ES	2107828 T	01-12-1997
			GR	3025194 T	27-02-1998
DE 4344646	A	29-06-1995	NONE		
WO 9701097	A	09-01-1997	US	5710009 A	20-01-1998
			AU	6387596 A	22-01-1997
			CA	2225210 A	09-01-1997
			EP	0840895 A	13-05-1998
			JP	10511776 T	10-11-1998

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.